

International Journal of Modern Physics E
© World Scientific Publishing Company

Quarkonia measurements with the central detectors of ALICE

Wolfgang Sommer, Christoph Blume, Frederick Kramer
*Institut für Kernphysik, University of Frankfurt,
Max-von-Laue Str. 1, Frankfurt, 60438, Germany
wolfgang.sommer@ikf.uni-frankfurt.de*

Jan Fiete Grosse-Oetringhaus
*Institut für Kernphysik, University of Münster,
Wilhelm-Klemm-Str. 9, Münster, 48149, Germany*

for the ALICE collaboration

Received (received date)

Revised (revised date)

A Large Ion Collider Experiment – ALICE will become operational with the startup of the Large Hadron Collider – LHC at the end of 2007. One focus of the physics program is the measurement of quarkonia in proton-proton and lead-lead collisions. Quarkonia states will be measured in two kinematic regions and channels: di-muonic decays will be measured in the forward region by the muon arm, the central part of the detector will measure di-electronic decays. The presented studies show the expected performance of the di-electron measurement in proton-proton and central lead-lead collisions.

1. Introduction

Already 1985, the suppression of quarkonia in heavy ion collisions relative to proton-proton collisions was suggested as a sign for deconfined matter that is produced in these collisions¹. Over the last two decades various experiments^{2 3} measured a significant suppression signal. Although strong indications for a J/Ψ suppression were found the theoretical picture remains unclear. At the LHC with the unprecedented center of mass energy of 5.5 TeV per nucleon in lead-lead collisions the deconfined phase is expected to be larger and longer lived. Thus one expects new insights into the nature of quarkonia suppression.

2. Quarkonia production at LHC energies

To estimate the quarkonia production cross section, predictions from the Color Evaporation Model (CEM)⁴ were used. The cross sections are calculated as a product of the $q\bar{q}$ cross section and a transition probability to the individual states. Possible feed down from higher mass states was also included. A comparison to measured cross sections at the highest available Tevatron energy was done and the

2 *W. Sommer, C. Blume, J.F. Grosse-Oetringhaus, F. Kramer*

Table 1. Expected cross sections for quarkonia at the LHC. The cross sections include the branching into di-electrons as well as feed down from higher states. All numbers are given in μb .

System	\sqrt{s}	J/Ψ	Ψ'	Υ	Υ'	Υ''
proton-proton	14 TeV	3.18	0.057	0.02	0.005	0.003
proton-proton	5.5 TeV	1.83	0.033	0.009	0.002	0.0013
lead-lead	5.5 TeV	48930	879	304	78.8	44.4

calculations were tuned to reproduce experimental data. The scaling from proton-proton to lead-lead collisions was done using the number of binary collisions, taking into account nuclear modifications employing the EKS98⁵ parametrization. The resulting cross sections are shown in Table 1.

3. The ALICE detector

ALICE will be the only dedicated heavy-ion experiment at the LHC. It was optimized to measure a large variety of heavy-ion observables at a centre of mass energy of 5.5 TeV per nucleon. Besides others the measurement of quarkonia is a key element of the physics program. Quarkonia decaying into di-muons will be measured in the muon arm covering forward rapidities ($2.5 > y > 4.0$)⁶. Di-electronic decays will be measured in the central barrel ($|y| < 0.9$) involving the following three detectors:

- **Inner Tracking System (ITS)**

The Inner Tracking system consists of three different types of silicon detectors, a silicon pixel detector (SPD), a silicon strip detector (SSD) and a silicon drift detector (SDD) arranged in six layers around the beam axis. The resolution of the combined ITS is $\sim 70 \mu\text{m}$ in $r\phi$ and $\sim 170 \mu\text{m}$ in z at $p_t=1 \text{ GeV}/c$ and $\sim 20 \mu\text{m}$ ($60 \mu\text{m}$) at $p_t=10 \text{ GeV}/c$. Thus it will enable the measurement of open charm and beauty decays via their displaced vertices.

- **Time Projection Chamber (TPC)**

With a diameter of $\sim 5 \text{ m}$, a length of $\sim 5 \text{ m}$ and a volume of more than 95 m^3 the ALICE TPC is the largest TPC ever built. It is the main tracking device with a momentum resolution of less than 2.5% for 4 GeV/c electron tracks. Besides that it will also serve as a particle identification detector using the specific energy loss of particles in the TPC gas (90% Ne, 10% CO₂).

- **Transition Radiation Detector (TRD)**

The Transition Radiation Detector was added to the experiment to improve the tracking capabilities and to discriminate electrons from the huge background of pions and other hadrons. The TRD will also serve as a fast

trigger device for high p_t electrons. It cylindrically surrounds the TPC with an inner diameter of ~ 6 m and an outer diameter of ~ 6.8 m and is segmented into 18 *super modules*. Each of these super modules is segmented in 5 *stacks* in z-direction with 6 layers of readout chambers in radial direction per stack. This gives a total of 540 read out chambers covering an area of 736 m^2 with 1.16 million readout channels.

4. Simulation technique

All presented studies were done within the ALICE offline software framework *ali-root*. The aim was to simulate the expected performance of the di-electron quarkonia measurement with realistic estimates on the quarkonia production ratio, background and detector response. During one ALICE running year we expect to record $2 \cdot 10^9$ proton-proton collisions and $2 \cdot 10^8$ central lead-lead collisions. The complete simulation and reconstruction of one lead-lead event ($dN/dy_{ch} = 3000$) takes about 5 hours on a modern computer. It is clear that this amount of events exceeds the available computing facilities by far. For this reason a fast simulation framework was developed. Instead of using the time consuming full detector simulation a look up table was created⁷, parameterizing the detector efficiency and resolution with respect to the kinematic variables p_t , θ and ϕ as well as for particle identification. Figure 1 shows an example of the parameterization. The look up table was created




Fig. 1. Response function for efficiencies in p_t (a), θ (b) and ϕ (c). The efficiency for high p_t electrons decreases with p_t since more straight tracks tend to stay in dead zones while curved tracks have a higher probability to re-enter the detector. The efficiencies for θ and ϕ reflect the detector geometry. In θ a decrease due to the central electrode is visible and the 18 super modules of the TRD are nicely visible as dips in the ϕ -efficiency.

by analysing 5000 simulated Hijing-events with embedded electrons.

To generate the input to the fast simulations a cocktail of particles was created. The output of this generator was evaluated using the detector parameterization and analysed afterwards.

4 *W. Sommer, C. Blume, J.F. Grosse-Oetringhaus, F. Kramer*

5. Results

5.1. Proton-proton collisions




Fig. 2. The invariant mass distribution as obtained by the described simulation of $2 \cdot 10^8$ minimum bias proton-proton events. To the left the complete spectrum is shown while the right plot focuses on the J/Ψ region. The J/Ψ peak is clearly visible with 360 entries and a Gaussian width of $30 \text{ MeV}/c^2$. The signal to background ratio is 9 and the significance is 18. Apart from the J/Ψ no other quarkonia state gives a significant signal. However, one should note that a trigger on di-electrons will significantly enhance the expected yields.

For the evaluation of the proton-proton measurement performance $2 \cdot 10^8$ minimum bias events were generated⁸. This represents 20% of the expected events to be recorded during one proton-proton run and enough statistics to extrapolate the results to full statistics. The events were generated using a cocktail of quarkonia simulated with the cross sections given in Table 1 and a proton-proton PYTHIA⁹ event simulating the background including open charm and beauty. A single electron p_t cut of $1 \text{ GeV}/c$ was applied. One finds a significant J/Ψ peak. Fitting this peak with a Gaussian gives 360 entries within the area of $\pm 1.5\sigma$ around the mean value and a width of $30 \text{ MeV}/c^2$. The like-sign distribution was used to estimate the background. Since like-sign pairs are produced with much less probability than unlike-sign pairs in proton-proton collisions the distribution had to be scaled to match the unlike-sign distribution. After scaling the distribution it was used for background estimations. Table 2 summarizes the obtained results.

5.2. Lead-lead collisions

Figure 3 shows the result for lead-lead collisions. $2 \cdot 10^8$ central events were simulated representing the expected statistics for one ALICE running year¹⁰. The event cocktail consisted of quarkonia signals which were simulated given the rates in Table 1. To account for electrons coming from semi-leptonic charm and beauty decays 100 $c\bar{c}$ and 4 $b\bar{b}$ per event were embedded. Pionic background was simulated using a parameterized HIJING generator tuned to the multiplicity of $dN_{ch}/dy=3000$.




Fig. 3. Invariant mass distribution as expected for $2 \cdot 10^8$ central lead-lead collisions. A single electron p_t cut of $1 \text{ GeV}/c$ was applied. The solid black line shows the unlike-sign distribution. The background was estimated using like-sign pairs (thin black line). The gray areas indicate the pure quarkonia distributions. The dashed lines show the various contributions from other sources to the distributions.




Fig. 4. The invariant mass spectrum in the J/Ψ (left) and Υ (right) mass region. The signal to background ratio for the J/Ψ is 1.2, for the $\Upsilon(1S)$ 1.1 and for the $\Upsilon(2S)$ 0.8. $\Psi(2S)$ and the $\Upsilon(3S)$ do not have a significant peak. A fit of a Gaussian to the observed peaks gives a width of $30 \text{ MeV}/c^2$ for the J/Ψ and $90 \text{ MeV}/c^2$ for the Υ .

There are significant signals for the J/Ψ , $\Upsilon(1S)$ and $\Upsilon(2S)$. $\Psi(2S)$ and $\Upsilon(3S)$ have only small contributions above background. The background was estimated using like-sign technique. After subtraction of the background and fitting the peaks with a Gaussian one obtains a width of $30 \text{ MeV}/c^2$ for the J/Ψ and $90 \text{ MeV}/c^2$ for the Υ , this resolution is sufficient to resolve the individual Υ states. All further results are summarized in Table 2.

6 *W. Sommer, C. Blume, J.F. Grosse-Oetringhaus, F. Kramer*

Table 2. Summary of the main results for minimum bias proton-proton and central lead-lead collisions. Values are only given for quarkonia states where significant signals are expected. The proton-proton results are scaled to $2 \cdot 10^9$ events (one ALICE running year). See text for detailed description.

System		J/Ψ	$\Upsilon(1S)$	$\Upsilon(2S)$
proton-proton $\sqrt{s}=14$ TeV $2 \cdot 10^9$ ev.	signals	1800	15	–
	S/B	9	–	–
	$S/\sqrt{S+B}$	40	–	–
lead-lead $\sqrt{s}=5.5$ TeV $2 \cdot 10^8$ ev.	signals	120.000	900	350
	S/B	1.2	1.1	0.35
	$S/\sqrt{S+B}$	245	21	8

6. Summary

The expected performance of the quarkonia measurement in the di-electron channel has been simulated under realistic assumptions. Within one year of running ALICE will accumulate enough statistics to measure J/Ψ s in proton-proton and J/Ψ s, $\Upsilon(1S)$ and $\Upsilon(2S)$ in lead-lead collisions with good signal to background ratio and significance. The invariant mass resolution is sufficient to resolve the individual Υ states.

Acknowledgements

This work was supported by the German Federal Ministry of Education and Research.

References

1. T. Matsui and H. Satz, *J/Ψ suppression by Quark-Gluon Plasma formation*, Phys. Lett. **B178:416**, (1986).
2. B. Alessandro *et al.* [NA50 Collaboration], Eur. Phys. J. C **39** (2005) 335.
3. A. Adare [PHENIX Collaboration], arXiv:nucl-ex/0611020.
4. V. D. Barger, W. Y. Keung and R. J. Phillips, *Phys. Lett.* **B91** (1980) 253.
5. K. J. Eskola, V. J. Kolhinen and C. A. Salgado, *Eur. Phys. J.* **C9** (1999) 61.
6. G. Martínez-García for the ALICE collaboration, Proceedings of Quark Matter 2006, to be published in Journal of Physics G: Nuclear and Particle Physics (2007)
7. J.F. Grosse-Oetringhaus, *Determination of the Physics Performance of the ALICE Central Barrel using a distributed GRID Computing Environment*, Diploma thesis, IKP Münster (2005).
8. F. Kramer, *Studie zur Messung von Quarkonia mit dem ALICE-TRD und Aufbau eines Teststandes für seine Ausleskammern*, Diploma thesis, IKF Frankfurt (2006).
9. www.thep.lu.se/~torbjorn/Pythia.html
10. B. Alessandro *et al.* [ALICE Collaboration], ALICE: Physics performance report, volume II, J. Phys. G **32** (2006) 1295.